

# Mirage mediation, uplifting and gaugino masses

**Hans Peter Nilles**

Physikalisches Institut, Universität Bonn, Germany

Based on work with K. Choi, A. Falkowski, M. Olechowski, S. Pokorski,  
hep-th/0411066, hep-th/0503216, hep-ph/0702146

O. Lebedev, Y. Mambrini, V. Loewen, M. Ratz, hep-th/0603047, hep-0612035

# Outline

- Basic questions: moduli stabilization and Susy breakdown: Fluxes and Gaugino Condensation
- A large and a little hierarchy
- Mirage Mediation
- Distinct pattern of soft terms
- Some remarks on fine tuning
- Explicit schemes KKLT and LNR
- Robust prediction for gaugino masses
- The Gaugino Code
- Conclusions and outlook

# Two Basic Questions

- how to obtain Susy breakdown at a small scale?
- stabilization of moduli?

# Two Basic Questions

- how to obtain Susy breakdown at a small scale?
- stabilization of moduli?

Are the masses of moduli connected to Susy breakdown?  
Relevant moduli are

- Dilaton ( $S$ )
- Kähler ( $T_i$ ) and complex structure moduli ( $Z_\alpha$ )

# Two Basic Questions

- how to obtain Susy breakdown at a small scale?
- stabilization of moduli?

Are the masses of moduli connected to Susy breakdown?  
Relevant moduli are

- Dilaton ( $S$ )
- Kähler ( $T_i$ ) and complex structure moduli ( $Z_\alpha$ )
- Other moduli are needed.
- They might come from Chern-Simons terms, additional matter fields, hidden sectors.....

# Fluxes and gaugino condensation

Is there a general pattern of the soft mass terms?

We always have (from **flux** and **gaugino condensate**)

$$W = \text{something} - \exp(-X)$$

where “**something**” is small and  $X$  is moderately large.

# Fluxes and gaugino condensation

Is there a general pattern of the soft mass terms?

We always have (from **flux** and **gaugino condensate**)

$$W = \text{something} - \exp(-X)$$

where “**something**” is small and  $X$  is moderately large.

In fact in this simple scheme

$$X \sim \log(M_{\text{Planck}}/m_{3/2})$$

providing a “**little**” hierarchy.

(Choi, Falkowski, HPN, Olechowski, Pokorski, 2004)

# Mixed Modulus Anomaly Mediation

The contribution from “Modulus Mediation” is therefore suppressed by the factor

$$X \sim \log(M_{\text{Planck}}/m_{3/2})$$

Numerically this factor is given by:  $X \sim 4\pi^2$ .



# Mixed Modulus Anomaly Mediation

The contribution from “Modulus Mediation” is therefore suppressed by the factor

$$X \sim \log(M_{\text{Planck}}/m_{3/2})$$

Numerically this factor is given by:  $X \sim 4\pi^2$ .

Thus the contribution due to “Anomaly Mediation” (suppressed by a loop factor) becomes competitive, leading to a Mixed Modulus-Anomaly-Mediation scheme.

For reasons that will be explained later we call this scheme

**MIRAGE MEDIATION**

(Loaiza, Martin, HPN, Ratz, 2005)

# The little hierarchy

$$m_X \sim \langle X \rangle m_{3/2} \sim \langle X \rangle^2 m_{\text{soft}}$$

is a generic signal of such a scheme

(Choi, Falkowski, HPN, Olechowski, 2005)

# The little hierarchy

$$m_X \sim \langle X \rangle m_{3/2} \sim \langle X \rangle^2 m_{\text{soft}}$$

is a generic signal of such a scheme

(Choi, Falkowski, HPN, Olechowski, 2005)

- moduli and gravitino are heavy
- gaugino mass spectrum is compressed

# The little hierarchy

$$m_X \sim \langle X \rangle m_{3/2} \sim \langle X \rangle^2 m_{\text{soft}}$$

is a generic signal of such a scheme

(Choi, Falkowski, HPN, Olechowski, 2005)

- moduli and gravitino are heavy
- gaugino mass spectrum is compressed
- such a situation occurs if for some reason the Susy breaking is “sequestered”

# Mirage Unification

Mirage Mediation provides a

- characteristic pattern of soft breaking terms.

# Mirage Unification

Mirage Mediation provides a

- characteristic pattern of soft breaking terms.

To see this, let us consider the gaugino masses

$$M_{1/2} = M_{\text{modulus}} + M_{\text{anomaly}}$$

as a sum of two contributions of comparable size.

# Mirage Unification

Mirage Mediation provides a

- characteristic pattern of soft breaking terms.


To see this, let us consider the gaugino masses

$$M_{1/2} = M_{\text{modulus}} + M_{\text{anomaly}}$$

as a sum of two contributions of comparable size.


- $M_{\text{anomaly}}$  is proportional to the  $\beta$  function,  
i.e. **negative** for the gluino, **positive** for the bino
- thus  $M_{\text{anomaly}}$  is non-universal below the GUT scale

# Evolution of couplings





# The Mirage Scale



(Lebedev, HPN, Ratz, 2005)

# The Mirage Scale (II)

The gaugino masses coincide

- above the GUT scale
- at the mirage scale

$$\mu_{\text{mirage}} = M_{\text{GUT}} \exp(-8\pi^2/\rho)$$

# The Mirage Scale (II)

The gaugino masses coincide

- above the GUT scale
- at the mirage scale


$$\mu_{\text{mirage}} = M_{\text{GUT}} \exp(-8\pi^2/\rho)$$

where  $\rho$  denotes the “ratio” of the contribution of **modulus** vs. **anomaly mediation**. We write the gaugino masses as

$$M_a = M_s(\rho + b_a g_a^2) = \frac{m_{3/2}}{16\pi^2}(\rho + b_a g_a^2)$$


and  $\rho \rightarrow 0$  corresponds to pure anomaly mediation.

# Constraints on the mixing parameter



(Löwen, HPN, Ratz, 2006)

# Constraints on $\rho$



(Löwen, HPN, Ratz, 2006)

# The “MSSM hierarchy problem”

The scheme predicts a rather high mass scale

- multi-TeV for the gravitino
- rather high mass for the LSP-Neutralino

# The “MSSM hierarchy problem”

The scheme predicts a rather high mass scale

- multi-TeV for the gravitino
- rather high mass for the LSP-Neutralino

Thus we might worry about a **fine-tuning** to obtain

- the mass of the weak scale around 100 GeV from

$$\frac{m_Z^2}{2} = -\mu^2 + \frac{m_{H_d}^2 - m_{H_u}^2 \tan^2 \beta}{\tan^2 \beta - 1},$$

and there are large corrections to  $m_{H_u}^2$  .....

(Choi, Jeong, Kobayashi, Okumura, 2005)

# The “MSSM hierarchy problem” solved?

The influence of the various soft terms is given by

$$m_Z^2 \simeq -1.8 \mu^2 + 5.9 M_3^2 - 0.4 M_2^2 - 1.2 m_{H_u}^2 + 0.9 m_{q_L^{(3)}}^2 + \\ + 0.7 m_{u_R^{(3)}}^2 - 0.6 A_t M_3 + 0.4 M_2 M_3 + \dots ,$$



# The “MSSM hierarchy problem” solved?

The influence of the various soft terms is given by

$$m_Z^2 \simeq -1.8 \mu^2 + 5.9 M_3^2 - 0.4 M_2^2 - 1.2 m_{H_u}^2 + 0.9 m_{q_L^{(3)}}^2 + \\ + 0.7 m_{u_R^{(3)}}^2 - 0.6 A_t M_3 + 0.4 M_2 M_3 + \dots ,$$

Mirage mediation improves the situation

- especially for **small  $\rho$**
- because of a **reduced gluino mass**

(Choi, Jeong, Kobayashi, Okumura, 2005)

- explicit model building required

(Lebedev, HPN, Ratz, 2005; Pierce, Thaler, 2006)

# Explicit schemes I

The different schemes depend on the mechanism of uplifting:

- **uplifting with anti D3 branes**

(Kachru, Kallosh, Linde, Trivedi, 2003)

- $\rho \sim 5$  in the original KKLT scenario leading to
- a **mirage scale** of approximately  $10^{11}$  GeV

# Explicit schemes I


The different schemes depend on the mechanism of uplifting:

- **uplifting with anti D3 branes**

(Kachru, Kallosh, Linde, Trivedi, 2003)


- $\rho \sim 5$  in the original KKLT scenario leading to
- a **mirage scale** of approximately  $10^{11}$  GeV
- This scheme leads to **pure mirage mediation**:
  - gaugino masses and
  - scalar masses
- **both meet at a common mirage scale**

# Constraints on $\rho$



(Löwen, HPN, Ratz, 2006)

# The Mirage Scale



(Lebedev, HPN, Ratz, 2005)

# Explicit schemes II

- uplifting via matter superpotentials

(Lebedev, HPN, Ratz, 2006)

- allows a continuous variation of  $\rho$
- leads to potentially new contributions to sfermion masses


# Explicit schemes II

- **uplifting via matter superpotentials**

(Lebedev, HPN, Ratz, 2006)

- allows a continuous variation of  $\rho$
- leads to potentially **new contributions** to sfermion masses
- **gaugino masses still meet at a mirage scale**
- **soft scalar masses might be dominated by modulus mediation**
- similar constraints on the mixing parameter


# Constraints on the mixing parameter



(V. Löwen, 2007)




# Constraints on the mixing parameter



(V. Löwen, 2007)

# Constraints on the mixing parameter



(V. Löwen, 2007)

# Explicit schemes III

- This “relaxed” mirage mediation is rather common for schemes with F-term uplifting  
(Gomez-Reino, Scrucca; Dudas, Papineau, Pokorski; Abe, Higaki, Kobayashi, Omura; Lebedev, Löwen, Mambrini, HPN, Ratz ,2006)
- although “pure” mirage mediation is possible as well

# Explicit schemes III

- This “relaxed” mirage mediation is rather common for schemes with F-term uplifting  
(Gomez-Reino, Scrucra; Dudas, Papineau, Pokorski; Abe, Higaki, Kobayashi, Omura; Lebedev, Löwen, Mambrini, HPN, Ratz ,2006)
- although “pure” mirage mediation is possible as well

## Main message

- predictions for gaugino masses are more robust than those for sfermion masses

mirage pattern for gaugino masses rather generic

# Obstacles to D-term uplifting

In supergravity we have the relation

$$D \sim \frac{F}{W}$$

which implies that KKLT AdS minimum cannot be uplifted via D-terms.

(Choi, Falkowski, HPN, Olechowski, 2005)

# Obstacles to D-term uplifting

In supergravity we have the relation

$$D \sim \frac{F}{W}$$

which implies that KKLT AdS minimum cannot be uplifted via D-terms.

(Choi, Falkowski, HPN, Olechowski, 2005)

Moreover in these schemes we have

$$F \sim m_{3/2} M_{\text{Planck}} \quad \text{and} \quad D \sim m_{3/2}^2.$$

So if  $m_{3/2} \ll M_{\text{Planck}}$  the D-terms are irrelevant.

(Choi, Jeong, 2006)

# The Gaugino Code

How can we test these ideas at the LHC?

Look for pattern of gaugino masses

Let us consider the

- low energy spectrum of the MSSM
- measured values of gauge coupling constants

$$g_1^2 : g_2^2 : g_3^2 \simeq 1 : 2 : 6$$

# The Gaugino Code

How can we test these ideas at the LHC?

Look for pattern of gaugino masses

Let us consider the

- low energy spectrum of the MSSM
- measured values of gauge coupling constants

$$g_1^2 : g_2^2 : g_3^2 \simeq 1 : 2 : 6$$

The evolution of gauge couplings would then lead to **unification** at a GUT-scale around  $10^{16}$  GeV



# The Gaugino Code

Observe that

- evolution of gaugino masses is tied to evolution of gauge couplings
- for MSSM  $M_a/g_a^2$  does not run (at one loop)

# The Gaugino Code

Observe that

- evolution of gaugino masses is tied to evolution of gauge couplings
- for MSSM  $M_a/g_a^2$  does not run (at one loop)
- if there are no strong threshold corrections at the high scale
- robust prediction for gaugino masses
- gaugino mass relations are the key to reveal the underlying scheme

3 characteristic patterns

(Choi, HPN, 2007)

# mSUGRA Pattern

Universal gaugino mass at the GUT scale

- mSUGRA pattern:

$$M_1 : M_2 : M_3 \simeq 1 : 2 : 6 \simeq g_1^2 : g_2^2 : g_3^2$$

as realized in popular schemes such as gravity-, modulus-, gauge- and gaugino-mediation

# mSUGRA Pattern

Universal gaugino mass at the GUT scale

- mSUGRA pattern:

$$M_1 : M_2 : M_3 \simeq 1 : 2 : 6 \simeq g_1^2 : g_2^2 : g_3^2$$

as realized in popular schemes such as gravity-, modulus-, gauge- and gaugino-mediation

This leads to

- LSP  $\chi_1^0$  predominantly Bino
- $M_{\text{gluino}}/m_{\chi_1^0} \simeq 6$

as a characteristic signature of these schemes.

# Anomaly Pattern

Gaugino masses below the GUT scale determined by the  $\beta$  functions

- anomaly pattern:

$$M_1 : M_2 : M_3 \simeq 3.3 : 1 : 9$$

at the TeV scale as the signal of anomaly mediation.

# Anomaly Pattern

Gaugino masses below the GUT scale determined by the  $\beta$  functions

- anomaly pattern:

$$M_1 : M_2 : M_3 \simeq 3.3 : 1 : 9$$

at the TeV scale as the signal of anomaly mediation.

For the gauginos, this implies

- LSP  $\chi_1^0$  predominantly Wino
- $M_{\text{gluino}}/m_{\chi_1^0} \simeq 9$

Pure anomaly mediation inconsistent, as sfermion masses are problematic in this scheme (tachyonic sleptons).

# Mirage Pattern

Mixed boundary conditions at the GUT scale characterized by the parameter  $\rho$  (the ratio of anomaly to modulus mediation).

- $M_1 : M_2 : M_3 \simeq 1 : 1.3 : 2.5$  for  $\rho \simeq 5$
- $M_1 : M_2 : M_3 \simeq 1 : 1 : 1$  for  $\rho \simeq 2$

# Mirage Pattern

Mixed boundary conditions at the GUT scale characterized by the parameter  $\rho$  (the ratio of anomaly to modulus mediation).

- $M_1 : M_2 : M_3 \simeq 1 : 1.3 : 2.5$  for  $\rho \simeq 5$
- $M_1 : M_2 : M_3 \simeq 1 : 1 : 1$  for  $\rho \simeq 2$

The mirage scheme leads to

- LSP  $\chi_1^0$  predominantly Bino
- $M_{\text{gluino}}/m_{\chi_1^0} < 6$
- a “compact” gaugino mass pattern.



# Conclusion

Mirage Mediation naturally appears in string theory models with background fluxes and gaugino condensation. It

- relieves cosmological problems of moduli and gravitino
- reduces the fine tuning of the weak scale
- gives a **consistent neutralino dark matter** candidate

# Conclusion

Mirage Mediation naturally appears in string theory models with background fluxes and gaugino condensation. It

- relieves cosmological problems of moduli and gravitino
- reduces the fine tuning of the weak scale
- gives a **consistent neutralino dark matter** candidate

## Mirage mediation

- avoids the problems of conventional schemes like anomaly and modulus mediation
- **is the correct way to implement anomaly mediation**
- gives a consistent picture with **very few parameters**

# Conclusion

The **source of Mirage Mediation** is the appearance of a small parameter

$$X^{-1} \sim \log(m_{3/2}/M_{\text{Planck}})$$

that leads to a (heavy) superpartner spectrum exhibiting

- a little hierarchy  $m_X \sim \langle X \rangle m_{3/2} \sim \langle X \rangle^2 m_{\text{soft}}$
- a rather heavy gravitino mass
- and an **unusual relation** between the gaugino masses.

# Conclusion

The **source of Mirage Mediation** is the appearance of a small parameter

$$X^{-1} \sim \log(m_{3/2}/M_{\text{Planck}})$$

that leads to a (heavy) superpartner spectrum exhibiting

- a little hierarchy  $m_X \sim \langle X \rangle m_{3/2} \sim \langle X \rangle^2 m_{\text{soft}}$
- a rather heavy gravitino mass
- and an **unusual relation** between the gaugino masses.

**Mirage Mediation provides a distinct pattern of soft terms that could be tested at the LHC!**

# Happy Birthday



# Happy Birthday



All the best for 60 + 60 years

# Bonn 2001



# Aspen 2004

