

## Exercises on Elementary Particle Physics

Prof. Dr. H.-P. Nilles

### 1. Dimensional Regularization of $\varphi^4$ Theory


Consider the Lagrangian of a real scalar field:

$$\mathcal{L} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - \frac{1}{2} m^2 \varphi^2 - \frac{\lambda}{4!} \varphi^4$$

The Feynman rules for  $\varphi^4$  theory are very simple:

$$\text{---} \sim \frac{i}{p^2 - m^2 + i\epsilon} \quad \text{---} \times \text{---} \sim -i\lambda$$

(a) At 1-loop order, the propagator will receive a correction:



To calculate this loop graph, we need one final *Feynman rule*: Integrate over the momentum of the particle running in the loop. If it is a fermion, pick up an extra factor of -1.

Using the Feynman rules given above, calculate the amplitude for the correction to the propagator, and call the resulting integral  $I(m)$ .

(b) The amplitude  $I(m)$  is divergent in  $d = 4$  dimensions. Generally, the integrals are “less divergent” when the dimension  $d$  decreases. Perform the integration in  $d$  dimensions:

$$I(d, m) = \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 - m^2 + i\epsilon}$$

We will later take  $d \rightarrow 4$ . Now, perform a Wick rotation:

- i. View  $k^0$  as a complex variable. Draw the complex  $k^0$ -plane. The integration is along the real axis. Mark the positions of the poles of the propagator.
- ii. Rotate the axes by  $90^\circ$ . The integration which ran from  $-\infty$  to  $+\infty$  now runs from  $-i\infty$  to  $+i\infty$ . Explain why this rotation does not change the value of the integral.

- iii. Change variables:  $k^0 = iq^0$ ,  $k^i = q^i$ . Perform the variable substitution in the integral. The  $q^0$  integration runs from  $-\infty$  to  $+\infty$ . Rename the variables  $q \rightarrow k$ .

Your final result should read

$$I(d, m) = \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2 + m^2}.$$

- (c) The integral only depends on the absolute value of  $k$ . Change to radial coordinates. You will need the area of the sphere in  $d$  dimensions which is given by

$$\text{Vol}(\partial B^d) = \frac{d\pi^{d/2}}{\Gamma(d/2 + 1)}.$$

Then substitute

$$x = \frac{m^2}{|k|^2 + m^2} \quad \leftrightarrow \quad |k|^2 + m^2 = \frac{m^2}{x} \quad \leftrightarrow \quad 1 - x = \frac{|k|^2}{|k|^2 + m^2}.$$

Your final result should read

$$I(d, m) = \frac{d(4\pi)^{-d/2}(m^2)^{d/2-1}}{2\Gamma(d/2 + 1)} \int_0^1 dx (1-x)^{d/2-1} x^{-d/2}.$$

- (d) Use the definition of Euler's beta function

$$B(\alpha, \beta) = \int_0^1 dx x^{\alpha-1} (1-x)^{\beta-1}$$

and the relations

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)}, \quad \Gamma(z + 1) = z\Gamma(z) \quad (1)$$

to arrive at the following expression for the 1-loop correction of the propagator:

$$I(d, m) = (4\pi)^{-d/2} (m^2)^{d/2-1} \Gamma(1 - d/2). \quad (2)$$

- (e) The gamma function  $\Gamma(z)$  has 1st order poles at negative integers and 0. The case  $d \rightarrow 4$  corresponds to  $1 - d/2 \rightarrow -1$ , so the result has a pole as expected. To isolate the pole, we have to expand  $\Gamma(1 - d/2)$  near  $d = 4$ . Define  $\epsilon = 4 - d$ . Using eq. (1), first show that

$$\Gamma(1 - d/2) = \frac{2}{\epsilon} \frac{\Gamma(1 + \epsilon/2)}{-1 + \epsilon/2}.$$

Now we can expand around  $\Gamma(1) \neq \pm\infty$ . Taylor expand the gamma function in the numerator and its denominator (geometric series). Discarding all terms linear and higher order in  $\epsilon$ , your final result should read

$$\Gamma(1 - d/2) = - \left( \frac{2}{\epsilon} + 1 + \Gamma'(1) \right). \quad (3)$$

- (f) The first 2 factors in eq. (2) also have to be expanded in powers of  $\epsilon$ , because if their expansion contains terms of order  $\epsilon$ , there is a non-vanishing contribution from their product with the pole of the gamma function. First, substitute  $\epsilon = 4 - d$  to obtain

$$(4\pi)^{-d/2} (m^2)^{d/2-1} = (4\pi)^{-2} m^2 (4\pi)^{\epsilon/2} (m^2)^{-\epsilon/2}.$$

A dimensionfull quantity cannot be expanded in a power series. We introduce the so-called *renormalization point*  $\mu$  which has the dimension of mass:

$$(4\pi)^{-d/2} (m^2)^{d/2-1} = (4\pi)^{-2} m^2 (\mu^2)^{-\epsilon/2} (4\pi)^{\epsilon/2} \left(\frac{m^2}{\mu^2}\right)^{-\epsilon/2} \quad (4)$$

Next, use the relation

$$x = \exp \log x = 1 + \log x + \frac{1}{2!}(\log x)^2 + \frac{1}{3!}(\log x)^3 + \dots$$


to expand the last 2 factors in eq. (4). Your final result should read

$$(4\pi)^{-2} m^2 (\mu^2)^{-\epsilon/2} \left(1 + \frac{\epsilon}{2} \log 4\pi - \frac{\epsilon}{2} \log \frac{m^2}{\mu^2} + \dots\right). \quad (5)$$

- (g) Substituting your results obtained in eqs. (3), (5) into the expression for the 1-loop correction of the propagator given in eq. (2), show the following result:

$$I(d, m) = -(4\pi)^{-2} m^2 (\mu^2)^{-\epsilon/2} \left(\frac{2}{\epsilon} + 1 + \Gamma'(1) - \log \frac{m^2}{4\pi\mu^2} + \mathcal{O}(\epsilon)\right)$$

- (h) At 1-loop, the 4-point-function will receive a correction which is of 2nd order in  $\lambda$ :



Calculate the amplitude for the correction to the 4-point-function, and call the resulting integral  $J(d, m, q)$ . Note that the resulting amplitude depends on the sum of incoming momenta  $q = p_1 + p_2$ . Do not forget to impose 4-momentum conservation at the vertices.

- (i) Prove the so-called *Feynman trick*:

$$\frac{1}{ab} = \int_0^1 dx \frac{1}{[xa + (1-x)b]^2}$$

Hint: Solve the integral by variable substitution  $y = (a - b)x + b$ .

(j) Use the Feynman trick to rewrite the correction to the amplitude:

$$J(d, m, q) = \int_0^1 dx \int \frac{d^d k}{(2\pi)^d} \frac{1}{[(k + xq)^2 - \Delta]^2}, \quad \Delta = -x(1-x)q^2 + m^2 \quad (6)$$

(k) It is clear that we may shift the integration variable  $k$  in eq. (6) to get

$$J(d, m, q) = \int_0^1 dx \int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 - \Delta)^2}.$$

Differentiating eq. (2) with respect to  $m^2$ , derive

$$\frac{d}{dm^2} I(d, m) = \int \frac{d^d k}{(2\pi)^d} \frac{i}{(k^2 - m^2 + i\epsilon)^2} = (4\pi)^{-d/2} (m^2)^{d/2-2} \Gamma(2 - d/2).$$

Use this result to obtain

$$J(d, m, q) = \int_0^1 dx i(4\pi)^{-d/2} \Delta^{d/2-2} \Gamma(2 - d/2).$$

(l) Proceeding analogously to (1e)-(1g), show the following result:

$$J(d, m, q) = i(4\pi)^{-2} (\mu)^{-\epsilon/2} \left( \frac{2}{\epsilon} + \Gamma'(1) - \log \frac{m^2}{4\pi\mu^2} - \int_0^1 dx \log \frac{\Delta}{m^2} \right)$$

Note: The integral can be solved explicitly, see e.g. *Gradshteyn, Ryzhik, "Table of Integrals, Series, and Products", p. 250, eq. 2.733.*

So far, we have **regularized** the divergent integrals, i.e. we have separated the infinite and finite parts. The procedure to get rid of the infinities using physical input data is called **renormalizing** the theory. We will deal with renormalization in one of the next exercise sheets.

## 2. Triplet Higgs and the $\rho$ parameter

On the last exercise sheet, we learned that the GWS theory makes use of a Higgs *doublet* to break the  $SU(2)_L \times U(1)_Y$  gauge symmetry. After diagonalization of the mass matrices with the help of the Weinberg angle  $\tan \theta_W = \frac{g'}{g}$ , the vector boson masses were found to be

$$M_{W^\pm}^2 = \frac{1}{4}v^2g^2, \quad M_Z^2 = \frac{1}{4}v^2(g^2 + g'^2), \quad M_A^2 = 0.$$

(a) Calculate the  $\rho$  parameter for the GWS model, where  $\rho$  is defined by

$$\rho := \frac{M_{W^\pm}^2}{M_Z^2 \cos^2 \theta_W}$$

Now consider again the Lagrangian from the last exercise sheet

$$\mathcal{L} = (D_\mu \Phi)^\dagger (D^\mu \Phi) + \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2 - \frac{1}{4} F_{\mu\nu}^i F^{i\mu\nu} - \frac{1}{4} G_{\mu\nu} G^{\mu\nu}$$

but  $\Phi$  is now a triplet  $(\phi^{++}, \phi^+, \phi^0)^T$  under  $SU(2)_L$  and the 3 dimensional generators of  $SU(2)_L$  are given by

$$T^1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad T^2 = \frac{i}{\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad T^3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

- (b) Calculate the hypercharge of  $\Phi$  and write down the covariant derivative.
- (c) Calculate the vacuum expectation value  $\langle \Phi^\dagger \Phi \rangle =: \frac{v^2}{2}$  of the Higgs field and define the shifted field.
- (d) Find the vector boson masses from the shifted Lagrangian. Calculate the  $\rho$  parameter for this model.

Experiment says  $\rho = 1$ . Which of the two models can you rule out?